

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number  
**WO 02/02548 A1**

- (51) International Patent Classification<sup>7</sup>: C07D 311/04, 215/20, A61K 31/47, A61P 25/16, A61K 31/435
- (74) Agent: ORION CORPORATION; Orion Pharma, Industrial Property Rights, P.O. Box 65, FIN-02101 Espoo (FI).
- (21) International Application Number: PCT/FI01/00613
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 28 June 2001 (28.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
20001593 3 July 2000 (03.07.2000) FI
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): ORION CORPORATION [FI/FI]; Orionintie 1, FIN-02200 Espoo (FI).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): PYSTYNEN, Jarmo [FI/FI]; Iivisniementie 6 A 5, FIN-02260 Espoo (FI). OVASKA, Martti [FI/FI]; Huhtakoukku 16 H 15, FIN-02340 Espoo (FI). VIDGREN, Jukka [FI/FI]; Aleksis Kivenkatu 74 B 29, FIN-00520 Helsinki (FI). LOTTA, Timo [FI/FI]; Ryytimaantie 3 H 28, FIN-01630 Vantaa (FI). YLIPERTTULA-IKONEN, Marjo [FI/FI]; Vuoritie 22, FIN-02340 Espoo (FI).
- Published:  
— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: COUMARIN DERIVATIVES WITH COMT INHIBITING ACTIVITY



(57) Abstract: Compounds of formula (I') wherein the two OH- substituents in the phenyl moiety are in a position ortho to one another and R1 in a position ortho to one of the hydroxy groups; and X, R<sub>1</sub> to R<sub>6</sub> are as defined in claims, exhibit COMT enzyme inhibiting activity so that they are useful as COMT inhibitors.

## COUMARIN DERIVATIVES WITH COMT INHIBITING ACTIVITY

### FIELD OF THE INVENTION

The present invention relates to coumarin derivatives and pharmaceutically acceptable salts and esters thereof. The invention further relates to pharmaceutical compositions thereof and to their use as inhibitors of catechol-O-methyltransferase (COMT) enzyme.

### BRIEF DESCRIPTION OF THE PRIOR ART

Compounds with COMT inhibiting activity are already known. For example, derivatives of catechols and isoflavones as COMT inhibitors have been disclosed i.a. in US-A-5 446 194, US-A-5 389 653 and, respectively, in US-A-3 973 608. COMT inhibitors are used i.a. in the treatment of Parkinson's disease. COMT-inhibitors have also indicated to be useful in the treatment of i.a. hypertension, heart failure and depression (cf. e.g. US-A-446 194 above) as well as inhibitors for the prevention of diabetic vascular dysfunctions (cf. WO-A-98 27973).

As to the known derivatives of coumarin, WO-A-93 16064 discloses coumarins with tyrosine kinase enzyme inhibiting activity to be used as antitumor agents. Furthermore, J.Mazur and T.Zawadowski (Acta Pol.Pharm., vol.54(5), 1997, p.371-374, see also Pol.J.Chem., vol.55(5), 1981, p.1151-5), D.Desai and R.H.Mehta (Indian J.Heterocycl.Chem., vol.6(3), 1997, p.241-244) and A.C.Jain et al. (Bull.Chem.Soc.Jpn, vol.52(4), 1979, p.1203-4) describe various coumarin derivatives with i.a. antibacteric, hypotensive, spasmolytic and/or antileukaemic activity.


### SUMMARY OF THE INVENTION

The object of the present invention is to provide further compounds with catechol-O-methyltransferase enzyme inhibiting activity.

The invention also provides compounds for the treatment of disorders or conditions wherein inhibition of COMT is indicated to be useful, as well as a use thereof for the manufacture of a medicament to be used as a COMT inhibiting agent. Furthermore, pharmaceutical compositions containing the present compounds are provided.

## DETAILED DESCRIPTION OF THE INVENTION

The invention thus provides compounds of the general formula I':



- 5 wherein the two OH- substituents in the phenyl moiety are in a position ortho to one another and R1 in a position ortho to one of the hydroxy groups;

X is O or NR<sub>7</sub>; wherein R<sub>7</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or -(C<sub>1</sub>-C<sub>6</sub>)alkyl-COOH;

R<sub>1</sub> is NO<sub>2</sub>, CN, CHO, CF<sub>3</sub> or (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-;

10

R<sub>2</sub> and R<sub>3</sub> are each selected independently from H, OH, halogen, NO<sub>2</sub>, SH, NH<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, OH-(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, SO<sub>2</sub>R<sub>10</sub>, -CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl, -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH and -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>; wherein

15

m is 0 or 1;

n is 0 or 1;

Y is -CO- or -CHOH-;

B is (C<sub>1</sub>-C<sub>6</sub>)alkylene or (C<sub>2</sub>-C<sub>6</sub>)alkenylene;


20

R<sub>8</sub> is phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclyl with one to four heteroatoms each selected independently from N, O and S, wherein the said phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclyl is optionally substituted with one to five substituents R<sub>9</sub> each selected independently from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub>, SH, NH<sub>2</sub>, CN, CHO, =O, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, CO-NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino-CO-, NHOH, CONHOH and SO<sub>2</sub>R<sub>10</sub>;

25

or R<sub>8</sub> is

3



or  $R_2$  and  $R_3$  form together  $-(CH_2)_r$ ; wherein  $r$  is 3, 4 or 5;


$R_4$  and  $R_5$  are independently H or  $(C_1-C_6)$ alkyl; or  $R_4$  and  $R_5$  form together  
5  $=O$ ,  $=S$  or  $=NR_{11}$ , wherein  $R_{11}$  is H or  $(C_1-C_6)$ alkyl;

$R_6$  is H,  $NO_2$ , CN, CHO, halogen,  $CF_3$  or  $(C_1-C_6)$ alkyl; and

$R_{10}$  is  $(C_1-C_6)$ alkyl,  $NH_2$ , OH or mono- or di $(C_1-C_6)$ alkylamino;

or pharmaceutically acceptable salts or pharmaceutically acceptable esters  
thereof.

10 As a subgroup of the compounds I' the invention provides new compounds of  
formula I,



15 wherein X,  $R_1$  to  $R_6$  are as defined above under the compounds of formula I',  
with the provisos that (a) when X is O,  $R_2$  is methyl,  $R_3$  is H,  $R_4$  and  $R_5$  form  
together  $=O$ ,  $R_6$  is H and the two hydroxy substituents are at 7- and 8-positions,  
20 then  $R_1$  is not CHO, (b) when X is O,  $R_2$  is H or methyl,  $R_3$  is H,  $R_4$  and  $R_5$  form  
together  $=O$ ,  $R_6$  is H and the two hydroxy substituents are at 6- and 7-positions,  
then  $R_1$  is not 8-CO-CH<sub>3</sub>, (c) when X is O,  $R_2$  and  $R_3$  are H,  $R_4$  and  $R_5$  are both  
methyl,  $R_6$  is H and the two hydroxy substituents are at 7- and 8-positions, then  $R_1$   
is not -CO-CH<sub>3</sub>,

25 or pharmaceutically acceptable salts or pharmaceutically acceptable esters  
thereof.

The compounds of formula I' and I exhibit COMT inhibiting activity and can  
thus be used as therapeutics for the treatment of diseases or conditions wherein

COMT inhibitors are indicated to be useful, e.g. for the treatment of Parkinson's disease.

The following subgroups (1) to (15) of compounds of formula I' or I taken alone or in any combination with each other are preferable,

5 (1) R<sub>1</sub> is NO<sub>2</sub>, CN, or CF<sub>3</sub>, e.g. NO<sub>2</sub> or CN, such as NO<sub>2</sub>;


(2) X is O;

(3) X is NR<sub>7</sub>; R<sub>7</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl, e.g. H;


10 (4) R<sub>2</sub> and R<sub>3</sub> are each selected independently from H, OH, halogen, NO<sub>2</sub>, NH<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, OH-(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, -CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, e.g. -CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH, e.g. -(B)<sub>m</sub>-COOH and -Y-B-COOH, and -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>; wherein n is 0 or 1; m is 0 or 1; Y is -CO- or -CHOH-, e.g. -CO-; B is (C<sub>1</sub>-C<sub>6</sub>)alkylene or (C<sub>2</sub>-C<sub>6</sub>)alkenylene, e.g. (C<sub>1</sub>-C<sub>6</sub>)alkylene; R<sub>8</sub> is phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-  
15 membered heterocyclidyl with one to four heteroatoms each selected independently from N, O and S, e.g. phenyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S (e.g. piperidyl, piperazinyl, morpholinyl, tetrazolyl, thienyl, furyl or pyridyl, e.g. tetrazolyl or pyridyl) each of which is  
20 optionally substituted with one to five, e.g. one to three, such as one or two, e.g. one, substituent(s) R<sub>9</sub> each selected independently from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub>, SH, NH<sub>2</sub>, CN, CHO, =O, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, -CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, -CO-NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino-CO-, -NHOH, -CONHOH and SO<sub>2</sub>R<sub>10</sub>, wherein R<sub>10</sub> is as  
25 defined above (e.g. (C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub> or OH), e.g. from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub>, NH<sub>2</sub>, CN, CHO, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, -CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, -CO-NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino-CO-, -NHOH and -CONHOH;  
30 such as from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl; e.g. from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub> and (C<sub>1</sub>-C<sub>6</sub>)alkyl;

5

(5)  $R_2$  and  $R_3$  are each selected independently from H,  $-\text{CO}-(\text{C}_1-\text{C}_9)\text{alkyl}$ ,  $-(\text{Y})_n-(\text{B})_m-\text{COOH}$ , e.g.  $-(\text{B})_m-\text{COOH}$  and  $-\text{Y}-\text{B}-\text{COOH}$ , and  $-(\text{Y})_n-(\text{B})_m-\text{R}_8$ ; wherein  $n$  is 0 or 1;  $m$  is 0 or 1;  $\text{Y}$  is  $-\text{CO}-$  or  $-\text{CHOH}-$ , e.g.  $-\text{CO}-$ ;  $\text{B}$  is  $(\text{C}_1-\text{C}_6)\text{alkylene}$  or  $(\text{C}_2-\text{C}_6)\text{alkenylene}$ , e.g.  $(\text{C}_1-\text{C}_6)\text{alkylene}$ ;  $\text{R}_8$  is phenyl, naphthyl,  $(\text{C}_3-\text{C}_7)\text{cycloalkyl}$  or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S, e.g. phenyl,  $(\text{C}_3-\text{C}_7)\text{cycloalkyl}$  or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S (e.g. piperidyl, piperazinyl, morpholinyl, tetrazolyl, thienyl, furyl or pyridyl, e.g. tetrazolyl or pyridyl) each of which is unsubstituted or substituted with one to five, e.g. one to three, such as one or two, e.g. one, substituent(s)  $\text{R}_9$  each selected independently from OH, halogen, COOH, 5-tetrazolyl, NHOH, CONHOH,  $\text{NO}_2$ , SH,  $\text{NH}_2$ , CN, CHO,  $=\text{O}$ ,  $(\text{C}_1-\text{C}_6)\text{alkyl}$ ,  $(\text{C}_1-\text{C}_6)\text{alkoxy}$  and halo- $(\text{C}_1-\text{C}_6)\text{alkyl}$ , e.g.  $\text{CF}_3$ , or  $\text{R}_8$  is



(6) one of  $R_2$  and  $R_3$  is selected from  $-\text{CO}-(\text{C}_1-\text{C}_9)\text{alkyl}$ ,  $-(\text{Y})_n-(\text{B})_m-\text{COOH}$ , e.g.  $-(\text{B})_m-\text{COOH}$  and  $-\text{Y}-\text{B}-\text{COOH}$ , and  $-(\text{Y})_n-(\text{B})_m-\text{R}_8$ ; wherein  $n$  is 0 or 1;  $m$  is 0 or 1;  $\text{Y}$  is  $-\text{CO}-$  or  $-\text{CHOH}-$ , e.g.  $-\text{CO}-$ ;  $\text{B}$  is  $(\text{C}_1-\text{C}_6)\text{alkylene}$  or  $(\text{C}_2-\text{C}_6)\text{alkenylene}$ , e.g.  $(\text{C}_1-\text{C}_6)\text{alkylene}$ ;  $\text{R}_8$  is phenyl, naphthyl,  $(\text{C}_3-\text{C}_7)\text{cycloalkyl}$  or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S, e.g. phenyl,  $(\text{C}_3-\text{C}_7)\text{cycloalkyl}$  or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S (e.g. piperidyl, piperazinyl, morpholinyl, tetrazolyl, thienyl, furyl or pyridyl, e.g. 5-tetrazolyl or pyridyl) each of which is unsubstituted or substituted with one to five, e.g. one to three, such as one or two, e.g. one, substituent(s)  $\text{R}_9$  each selected independently from OH, halogen, COOH, 5-tetrazolyl, NHOH, CONHOH,  $\text{NO}_2$ , SH,  $\text{NH}_2$ , CN, CHO,  $=\text{O}$ ,  $(\text{C}_1-\text{C}_6)\text{alkyl}$ ,  $(\text{C}_1-\text{C}_6)\text{alkoxy}$  and halo- $(\text{C}_1-\text{C}_6)\text{alkyl}$  (e.g.  $\text{CF}_3$ ), e.g. from COOH, 5-tetrazolyl, NHOH and CONHOH, e.g. from COOH and 5-tetrazolyl; or  $\text{R}_8$  is



- (7) one of  $R_2$  and  $R_3$  is selected from H, OH, halogen,  $\text{NO}_2$ ,  $\text{NH}_2$ , ( $\text{C}_1$ - $\text{C}_6$ )alkyl, ( $\text{C}_2$ - $\text{C}_6$ )alkenyl, ( $\text{C}_1$ - $\text{C}_6$ )alkoxy, OH-( $\text{C}_1$ - $\text{C}_6$ )alkyl, halo-( $\text{C}_1$ - $\text{C}_6$ )alkyl, mono- or di( $\text{C}_1$ - $\text{C}_6$ )alkylamino, -CO-( $\text{C}_1$ - $\text{C}_9$ )alkyl and  $-(Y)_n$ -( $B$ ) $_m$ - $R_8$ ; e.g. one of  $R_2$  and  $R_3$  is  $-(Y)_n$ -( $B$ ) $_m$ - $R_8$ , wherein  $n$  is 0 or 1;  $m$  is 0 or 1;  $Y$  is -CO- or -CHOH-, e.g. -CO-;  $B$  is ( $\text{C}_1$ - $\text{C}_6$ )alkylene or ( $\text{C}_2$ - $\text{C}_6$ )alkenylene, e.g. ( $\text{C}_1$ - $\text{C}_6$ )alkylene;  $R_8$  is phenyl, naphthyl, ( $\text{C}_3$ - $\text{C}_7$ )cycloalkyl or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S, e.g. phenyl, ( $\text{C}_3$ - $\text{C}_7$ )cycloalkyl or 5- or 6-membered heterocyclic ring with one to four heteroatoms each selected independently from N, O and S (e.g. piperidyl, piperazinyl, morpholinyl, tetrazolyl, thienyl, furyl or pyridyl, suitably 5-tetrazolyl or pyridyl) each of which is unsubstituted or substituted with one to five, e.g. one to three, such as one or two, e.g. one, substituent(s)  $R_9$  each selected independently from OH, halogen,  $\text{COOH}$ , 5-tetrazolyl,  $\text{NO}_2$ , SH,  $\text{NH}_2$ , CN, CHO, =O, ( $\text{C}_1$ - $\text{C}_6$ )alkyl, ( $\text{C}_1$ - $\text{C}_6$ )alkoxy and halo-( $\text{C}_1$ - $\text{C}_6$ )alkyl (e.g.  $\text{CF}_3$ ), e.g. from OH, halogen,  $\text{NO}_2$ , CN, CHO, ( $\text{C}_1$ - $\text{C}_6$ )alkyl, ( $\text{C}_1$ - $\text{C}_6$ )alkoxy and halo-( $\text{C}_1$ - $\text{C}_6$ )alkyl (e.g.  $\text{CF}_3$ ); such as from halogen,  $\text{NO}_2$ , ( $\text{C}_1$ - $\text{C}_6$ )alkyl and ( $\text{C}_1$ - $\text{C}_6$ )alkoxy;
- (8)  $R_2$  is selected from (7) above;
- (9) one of  $R_2$  and  $R_3$ , e.g.  $R_3$ , is -CO-( $\text{C}_1$ - $\text{C}_9$ )alkyl,  $-(Y)_n$ -( $B$ ) $_m$ - $\text{COOH}$ , e.g. -( $B$ ) $_m$ - $\text{COOH}$  or -Y-B- $\text{COOH}$ , or  $-(Y)_n$ -( $B$ ) $_m$ - $R_8$ , such as -( $B$ ) $_m$ - $\text{COOH}$ , -Y-B- $\text{COOH}$  or  $-(Y)_n$ -( $B$ ) $_m$ - $R_8$ , wherein  $n$  is 0 or 1;  $Y$  is CO;  $m$  is 0 or 1,  $B$  is ( $\text{C}_1$ - $\text{C}_6$ )alkylene, e.g. - $\text{CH}_2$ -; and  $R_8$  is phenyl unsubstituted or substituted with  $\text{COOH}$  or tetrazol; or  $R_8$  is tetrazol, e.g. 5-tetrazol, or pyridyl.
- (10)  $Y$  is CO;
- (11)  $B$  is ( $\text{C}_1$ - $\text{C}_6$ )alkylene;

7


(12) one of  $R_2$  and  $R_3$  is as defined in any of (4) to (7) and/or (9) above and the other of  $R_2$  and  $R_3$  is selected independently from H, OH, halogen,  $\text{NO}_2$ ,  $\text{NH}_2$ ,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_2\text{-C}_6)\text{alkenyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , OH- $(\text{C}_1\text{-C}_6)\text{alkyl}$ , halo- $(\text{C}_1\text{-C}_6)\text{alkyl}$  and mono- or di $(\text{C}_1\text{-C}_6)\text{alkylamino}$ ; e.g. H, OH,  $(\text{C}_1\text{-C}_6)\text{alkyl}$  and  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ ; e.g. H, OH and  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ;

(13)  $R_2$  and  $R_3$  form together  $-(\text{CH}_2)_r$ ,  $r$  is 3, 4 or 5, e.g. 3 or 4;

(14)  $R_6$  is H,  $\text{NO}_2$ , CHO, halogen,  $\text{CF}_3$  or  $(\text{C}_1\text{-C}_6)\text{alkyl}$  (e.g. methyl); such as H,  $\text{NO}_2$ , CHO or  $(\text{C}_1\text{-C}_6)\text{alkyl}$  (e.g. methyl); e.g. H; and/or


(15)  $R_4$  and  $R_5$  are independently H or  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ; e.g.  $R_4$  and  $R_5$  are both methyl; or  $R_4$  and  $R_5$  form together  $=\text{O}$ ,  $=\text{S}$  or  $=\text{NH}$ ; e.g.  $=\text{O}$  or  $=\text{NH}$ ; such as  $=\text{O}$ .

A subgroup of the compounds of formula I' or I are the compounds of formula Ia,



wherein X,  $R_1$  to  $R_5$  are as defined above.

A further subgroup of the compounds of formula I' or I are the compounds of formula Ib,



wherein X,  $R_1$  to  $R_5$  are as defined above.



In a further subgroup of compounds of formula I', I, Ia or Ib, R<sub>1</sub> is NO<sub>2</sub> or CN, e.g. NO<sub>2</sub>, and one of R<sub>2</sub> and R<sub>3</sub>, e.g. R<sub>3</sub>, is -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH, e.g. -(B)<sub>m</sub>-COOH or -Y-B-COOH, or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>. In another subgroup of compounds I', I, Ia or Ib one of R<sub>2</sub> and R<sub>3</sub>, e.g. R<sub>3</sub>, is -(B)<sub>m</sub>-COOH, -Y-B-COOH or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>, n is 0 or 1, Y is CO; m is 0 or 1; B is (C<sub>1</sub>-C<sub>6</sub>)alkylene; R<sub>8</sub> is 5-tetrazolyl, or R<sub>8</sub> is phenyl or pyridyl unsubstituted or substituted with one or two, e.g. one, substituent(s) selected independently from OH, halogen, NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, COOH and 5-tetrazolyl. In another subgroup of compounds I', I, Ia or Ib one of R<sub>2</sub> and R<sub>3</sub>, e.g. R<sub>3</sub>, is -CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl, -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH, e.g. -(B)<sub>m</sub>-COOH or -Y-B-COOH, or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>.

In another subgroup of the compounds I', I, Ia or Ib, one of R<sub>2</sub> and R<sub>3</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>, wherein n, m Y and R<sub>8</sub> are as defined above; and the other of R<sub>2</sub> and R<sub>3</sub> is -CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl, -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH, e.g. -(B)<sub>m</sub>-COOH or -Y-B-COOH, or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub> as defined above. For example one of R<sub>2</sub> and R<sub>3</sub>, e.g. R<sub>2</sub>, is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and the other of R<sub>2</sub> and R<sub>3</sub>, e.g. R<sub>3</sub>, is -(B)<sub>m</sub>-COOH, -Y-B-COOH or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub> as defined above, e.g. -(B)<sub>m</sub>-COOH or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>, e.g. (C<sub>1</sub>-C<sub>6</sub>)alkylene-COOH or -(Y)<sub>n</sub>-R<sub>8</sub>, Y is CO, R<sub>8</sub> is phenyl or pyridyl unsubstituted or substituted with one or two, e.g. one, substituent(s) selected independently from OH, halogen, NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, COOH and 5-tetrazolyl, e.g. from OH, halogen, NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkoxy; or R<sub>8</sub> is tetrazolyl, e.g. 5-tetrazolyl.

In a further subgroup of the compounds I', I, Ia or Ib, R<sub>2</sub> and R<sub>3</sub> form together -(CH<sub>2</sub>)<sub>r</sub>, r is 3 or 4, e.g. 4. Furthermore, preferably in the compounds of formula I', I, Ia or Ib, R<sub>4</sub> and R<sub>5</sub> form together =O or =NH, e.g. =O, or R<sub>4</sub> and R<sub>5</sub> are both (C<sub>1</sub>-C<sub>6</sub>)alkyl, e.g. methyl.

In a preferred subgroup of the compounds of formula I' or I, the two hydroxy substituents at the phenyl ring are in 6- and 7-positions. Preferably, one of R<sub>2</sub> and R<sub>3</sub> is not H.

The compounds of formula I' and the subgroups I, Ia and Ib, as well as the pharmaceutically acceptable salts and the pharmaceutically acceptable esters thereof, are referred to below as the compounds of the invention, unless otherwise indicated.

The compounds of the invention may have chiral carbon atom(s) in their structure. The invention includes within its scope all the possible stereoisomers of

the compounds I, including geometric isomers, e.g. Z and E isomers (cis and trans isomers), and optical isomers, e.g. diastereomers and enantiomers. Furthermore, the invention includes in its scope both the individual isomers and any mixtures thereof, e.g. racemic mixtures. The individual isomers may be obtained using the  
5 corresponding isomeric forms of the starting material or they may be separated after the preparation of the end compound according to conventional separation methods. For the separation of i.a. optical isomers, e.g. enantiomers, from the mixture thereof the conventional resolution methods, e.g. fractional crystallisation, may be used.

10 Physiologically acceptable salts may be prepared by known methods. The pharmaceutically acceptable salts are the usual organic and inorganic salts in the art. Furthermore, the COOH-, OH- and/or amino-functionality, such as COOH- and/or OH-functionality, e.g. COOH- functionality, when present in the compounds of the invention, can be converted to a pharmaceutically acceptable ester or,  
15 respectively, a pharmaceutically acceptable amide in a manner known in the art using a pharmaceutically acceptable acid or, respectively, a pharmaceutically acceptable alcohol known from the literature. Examples of such pharmaceutically acceptable acids and alcohols are e.g. aliphatic (e.g. C<sub>1</sub>-C<sub>9</sub>, such as C<sub>1</sub>-C<sub>6</sub>) acids and alcohols, or aromatic acids and alcohols, which are conventional in the field of  
20 pharmaceuticals and which retain the pharmacological properties of the free form.

Terms employed herein have the following meanings: A halogen or halo refers to fluorine, chlorine, bromine or iodine. The term (C<sub>1</sub>-C<sub>6</sub>)alkyl as employed herein as such or as part of another group includes both straight and branched  
25 chain radicals of up to 6 carbon atoms, preferably of 1, 2, 3 or 4 carbon atoms. In CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl, the alkyl moiety includes both straight and branched chain radicals of up to 9 carbon atoms, preferably of up to 6 carbon atoms, e.g. 1, 2, 3 or 4 carbon atoms. The term (C<sub>1</sub>-C<sub>6</sub>)alkoxy as such or as part of another group refers to -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein (C<sub>1</sub>-C<sub>6</sub>)alkyl is as defined above. The term (C<sub>2</sub>-C<sub>6</sub>)alkenyl includes both straight and branched chain radicals of up to 6 carbon  
30 atoms, preferably of 2, 3 or 4 carbon atoms, containing double bond(s), e.g. one double bond. The term halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl refers to (C<sub>1</sub>-C<sub>6</sub>)alkyl radical, as defined above, that is substituted by one or more halo radicals as defined above, e.g. trifluoromethyl, difluoromethyl etc. The term (C<sub>1</sub>-C<sub>6</sub>)alkylene refers to a straight or branched, saturated hydrocarbon chain divalent radical, e.g. methylene, ethylene,  
35 propylene, butylene and the like. The term (C<sub>2</sub>-C<sub>6</sub>)alkenylene refers to a straight or branched, unsaturated hydrocarbon chain divalent radical, wherein the

10


unsaturation is present as one or more, e.g. one, double bond(s), e.g. vinylene, propenylene, butenylene etc. The term (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl refers to a monocyclic 3- to 7-membered saturated carbocyclic ring, i.e. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl ring. The 5- or 6-membered heterocyclyl with one to four heteroatoms selected independently from N, O and S means a monocyclic, partially or fully saturated, or aromatic hetero ring system. Examples of such heterocyclyls include piperidiny, piperaziny, morpholiny, pyrroly, tetrahydropyridyl, dihydropyridyl, pyridyl, pyraziny, pyrimidiny, pyridatsiny, thienyl, furyl, thiazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl etc., such as tetrazolyl, e.g. 5-tetrazolyl, thienyl or pyridyl.

In case of di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, the (C<sub>1</sub>-C<sub>6</sub>)alkyl chains can be identical or different.

It is evident to a skilled person that in the compounds I', I, Ia or Ib the nature of the optional substituent(s) R<sub>9</sub> and the maximal possible number thereof in a ring R<sub>8</sub> depend on the nature of the ring R<sub>8</sub>. E.g. the option =O as R<sub>9</sub> is possible only for (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or saturated or partially saturated heterocyclic rings as R<sub>8</sub>, wherein a double bond can be formed between the ring atom of R<sub>8</sub> and the said oxygen atom.

The compounds of the invention can be prepared by a variety of synthetic routes analogously or according to the methods known in the literature using suitable starting materials.


In general, the compounds of formula I', I, Ia or Ib, wherein R<sub>4</sub> and R<sub>5</sub> form together =O (compound Ic), can be prepared e.g. analogously or according to scheme 1:



wherein X, R<sub>1</sub> to R<sub>3</sub> and R<sub>6</sub> are as defined above, each R' is independently H or a conventional protecting group for hydroxy, e.g. (C<sub>1</sub>-C<sub>6</sub>)alkyl, and R is (C<sub>1</sub>-C<sub>6</sub>)alkyl, e.g. methyl or ethyl.

- 5 Accordingly, reaction of scheme 1 corresponds to the known von Pechman reaction. A compound II is reacted with a compound III in acidic reaction conditions, e.g. in HCl/alcohol or 75 % H<sub>2</sub>SO<sub>4</sub>, in a temperature between 0 °C and a room temperature, to obtain a compound Ic' which is then deprotected and a substituent R<sub>1</sub>, e.g. NO<sub>2</sub>, is introduced in a conventional manner to obtain a  
10 compound Ic.

The compounds I', I, Ia or Ib, wherein R<sub>4</sub> and R<sub>5</sub> form together =O (compound Ic), can further be prepared e.g. analogously or according to scheme 2:




- 15 wherein X, R<sub>1</sub> to R<sub>3</sub> and R<sub>6</sub> are as defined above, each R' is independently H or a conventional protecting group for hydroxy, e.g. (C<sub>1</sub>-C<sub>6</sub>)alkyl, and L is a conventional leaving group, e.g. OH, O-(C<sub>1</sub>-C<sub>6</sub>)alkyl or halogen.

- The reaction of scheme 2 is analogous to the known Knövenagel-  
condensation. In general, a compound IV is condensed in a suitable solvent, e.g.  
20 an alcohol, DMF, an alcoholic DMF, or THF, with a compound V in the presence of a base, e.g. an inorganic fluoride compound or an organic amine, such as piperidine, at an elevated temperature to obtain a compound Ic' which is deprotected to compound Ic as defined above.

12

More specifically, e.g. compounds I', I, Ia or Ib, wherein R<sub>4</sub> and R<sub>5</sub> form together =O, R<sub>2</sub> is H and R<sub>3</sub> is -(B)<sub>m</sub>-COOH, (compound Id), can also be prepared e.g. analogously or according to scheme 3:




5

wherein X and R<sub>1</sub> and R<sub>6</sub> are as defined above, each R' is independently H or a conventional protecting group for hydroxy, e.g. (C<sub>1</sub>-C<sub>6</sub>)alkyl, and R is (C<sub>1</sub>-C<sub>6</sub>)alkyl, e.g. methyl or ethyl.

The reaction of scheme 3 is analogous to the known Stobbe condensation. Accordingly, a compound VI is reacted with a compound VII in a suitable solvent, e.g. an alcohol, in the presence of a strong base, such as (C<sub>1</sub>-C<sub>6</sub>)alkyl-OMe, wherein Me is a metal ion, e.g. an alkaline metal ion, such as Na, in an elevated temperature to obtain a compound VIII. The compound VIII is cyclized and demethylated in a manner known in the art using a known demethylation reagent, e.g. borontribromide, at a cooled temperature, e.g. (-20) °C, and then R<sub>1</sub>, e.g. a nitro group, is introduced in a conventional manner to the resulted compound Id' to obtain compound Id as defined above.

A further method for preparing compounds I, wherein R<sub>4</sub> and R<sub>5</sub> form together =O, R<sub>1</sub> is NO<sub>2</sub> and X is NR<sub>7</sub>, wherein R<sub>7</sub> is as defined above, (compound Ie) is illustrated in scheme 4:


20



wherein R<sub>2</sub>, R<sub>3</sub> and R<sub>6</sub> are as defined above, R' is a conventional protecting group for hydroxy, e.g. (C<sub>1</sub>-C<sub>6</sub>)alkyl, and R is (C<sub>1</sub>-C<sub>6</sub>)alkyl.

- Accordingly, compound IX is reacted with compound III, in the presence or  
 5 absence of a suitable solvent, at elevated temperature, e.g. at about 160°C, at a period of time. Then, preferably without the isolation of the reaction product, an acid is added to the reaction mixture at a temperature of 50 - 100°C to obtain compound X which is optionally alkylated in the next step with Hal-R<sub>7</sub>, wherein Hal is halogen and R<sub>7</sub> is (C<sub>1</sub>-C<sub>6</sub>)alkyl. The compound X, or optionally XI, is  
 10 deprotected, e.g. demethylated, and then nitrated in a conventional manner to obtain the end compound 1e as defined above.

- Compounds I, wherein R<sub>4</sub> and R<sub>5</sub> are both (C<sub>1</sub>-C<sub>6</sub>)alkyl, e.g. methyl, and X is O, (compound If) can be prepared e.g. analogously or according to the method described by Cook et al. in J. Org. Chem., vol.30, 1965, p.4114, illustrated in  
 15 scheme 5:



wherein R<sub>1</sub> to R<sub>3</sub> and R<sub>6</sub> are as defined above and R' is a conventional protecting group for hydroxy, e.g. Me<sub>3</sub>Si.

Thus, compound XIII is reacted with MeMgCl in a suitable solvent, e.g. THF, and then deprotected with an acid to obtain the end compound If as defined above.

The starting materials II, III, IV, V, VI, VII, IX and XIII are commercially available or can be prepared via a variety of known synthetic routes known in the literature or described above.

It is obvious to a skilled person that, in the above reactions, any starting material or intermediate can be protected, if necessary, in a manner well known in the chemical field. Any protected functionality is subsequently deprotected in a usual manner.

Furthermore, in the above reaction schemes a substituent R<sub>1</sub> to R<sub>6</sub> and/or R<sub>7</sub> in the intermediates and/or end compounds of the invention may further be converted to another functionality of the invention, if desired, in a manner known to a skilled person.

It should be noted that the above described synthetic routes are meant to illustrate the preparation of the compounds of the invention and the preparation is by no means limited thereto, i.e. other synthetic methods which are within the general knowledge of a skilled person are also possible.

The compounds of the invention may be converted, if desired, into their pharmaceutically acceptable salt or ester form using methods well known in the art.

As already mentioned hereinbefore, the compounds of the invention show interesting pharmacological properties, namely they exhibit catechol-O-methyltransferase (COMT) enzyme inhibiting activity. The said activity of the compounds of the invention is demonstrated with the pharmacological tests presented below.

#### EXPERIMENT I: Determination of COMT activity (IC<sub>50</sub>)

The determination of IC<sub>50</sub> was performed by measuring the COMT activity in a test sample which contained S-COMT enzyme (about 30 nM), 3 mM dopamine (as the substrate of COMT), 5 mM magnesium chloride, 0.05 mM S-adenosyl-L-methionine (AdoMet) and a test compound of the invention at various concentrations in 0.1 M phosphate buffer, pH 7.4, at 37°C.

15

The reaction in the test sample was initiated by adding the dopamine substrate to the sample mixture and, after incubation for 15 min at 37°C, the reaction was stopped with 4 M perchloric acid and stabilized further 10 min in ice bath. Thereafter the precipitated proteins were removed by centrifugation (4000 x G for 10 min). The activity of COMT enzyme was measured by determining the concentration of the reaction products, 3-methyldopamine and 4-methyldopamine, by HPLC. The results were calibrated with 3-methyldopamine standards. See also T.Lotta et al., Biochemistry, vol.34(13), 1995, p.4204, T.Lotta et al. The IC<sub>50</sub> value is the concentration of the test compound which causes a 50 % decrease in COMT activity. The results are shown in table 1.

Table 1.

The compound of example no.	IC <sub>50</sub> (nM)
Example 6(b)	30
Example 7	30
Example 8	50
Example 9(c)	20
Example 10(d)	35
Example 11(c)	10
Example 12(b)	10
Example 13(c)	60

Particularly, the compounds of the invention have preferable properties as therapeutics. They can be used for the treatment of diseases or conditions wherein COMT inhibitors are indicated to be useful, i.a. in the treatment of Parkinson's disease for the potentiation of levodopa (+ DDC) therapy.

The compounds of the invention may be administered enterally, topically or parenterally.



The compounds of the invention may be formulated alone or together with one or more active agents and/or together with a pharmaceutically acceptable excipient in different pharmaceutical unit dosage forms, e.g. tablets, capsules, solutions, emulsions and powders etc., depending on the route of administration, using conventional techniques. The pharmaceutically acceptable excipient can be selected from those conventionally used in the field of pharmaceuticals noticing the chosen route of administration.

The amount of the active ingredient varies from 0.01 to 100 weight-% depending on i.a. the type of the dosage form.

The specific dose level of the compounds of the invention depends, of course, on several factors such as the compound to be administered, the species, age and the sex of the subject to be treated, the condition to be treated and on the route and method of administration. For example, the compounds of the invention may administered from 0.5  $\mu\text{g/kg}$  to 100 mg/kg per day for an adult male.

The present invention also provides a compound of the invention or an ester or salt thereof for use in a method of treatment of human or animal body.

The present invention further provides a compound of the invention or an ester or salt thereof, as well as a pharmaceutical composition thereof, for use as a COMT inhibitor, i.a. for the treatment of diseases and conditions where inhibition of COMT enzyme is useful, e.g. for the treatment of Parkinson's disease. The use of the compounds of the invention for the manufacture of a medicament to be used for the above indications is also provided. The invention further relates to a method for the treatment of above indicated conditions or diseases, by administering to a subject in need of such treatment an effective amount of the compound of the invention or a pharmaceutically acceptable ester or salt thereof.

The present invention will be explained in more detail by the following examples. The examples are meant only for illustrating purposes and does not limit the scope of the invention which is defined in claims.

## PREPARATION EXAMPLE 1

## (1a) 2-Hydroxy-3,4-dimethoxy-5-nitro-benzaldehyde

To a solution of 3,4-dimethoxy-2-hydroxybenzaldehyde (0.5 g) in acetic acid (10 ml) was added fuming nitric acid (0.12 ml). The reaction mixture was poured  
5 into ice water, the product extracted into ethyl acetate, washed with water, dried and evaporated. The product was triturated with water and filtered. Yield: 0.15 g.

NMR (90 MHz): 3.82 (s, 3H, CH<sub>3</sub>O), 4.00 (s, 3H, CH<sub>3</sub>O), 8.02 (s, 1H, ArH), 10.20 (s, 1H, CHO).

## EXAMPLE 1:

## 10 (a) 7,8-Dimethoxy-6-nitro-3-(4-nitro-phenyl)-chromen-2-one

A solution of 2-hydroxy-3,4-dimethoxy-5-nitro-benzaldehyde obtained from Preparation Example (1a) (0.46 g), 4-nitrophenyl acetic acid (0.39 g) and triethylamine (0.84 ml) in 1,2-dichloroethane (30 ml) was treated with phenyl  
15 dichlorophosphate (0.44 ml) at 0° C. The mixture was refluxed for 5 hours and cooled. The cool reaction mixture was diluted with methylene chloride, washed successively with 2.5 M NaOH, 2 M HCl and water, dried and evaporated. The product was finally triturated with pethroleum ether and filtered. Yield: 0.52 g, melting point 219-223°C.

NMR (90 MHz): 4.08 (s, 3H, CH<sub>3</sub>O), 4.10 (s, 3H, CH<sub>3</sub>O), 7.98-8.5 (m, 6 H).

## 20 (b) 7,8-Dihydroxy-6-nitro-3-(4-nitro-phenyl)-chromen-2-one

The product from the previous step (a) (0.5 g) was refluxed in concentrated hydrogen bromide (10 ml) under nitrogen for 5 hours. The product was worked up in the usual way and recrystallized from methanol. Yield: 0.18 g, melting point 270-280°C. NMR (90 MHz): 7.92-8.04 (m, 3H), 8.2-8.45 (m, 3 H).

## 25 EXAMPLE 2:

## (a) 7,8-Dimethoxy-6-nitro-2-oxo-2H-chromene-3-carboxylic acid ethyl ester

A solution of 3,4-dimethoxy-2-hydroxy-5-nitrobenzaldehyde obtained from Preparation Example (1a) (0.46 g), diethylmalonate (0.61 ml) and piperidine (two drops) was refluxed overnight. The product crystallized on cooling. Yield: 0.30 g.

18

NMR (400 MHz): 1.32 (t, 3H, CH<sub>3</sub>CH<sub>2</sub>O, J = 7.5 Hz), 4.00 (s, 3H, MeO), 4.08 (s, 3H, MeO), 4.32 (q, 2H, CH<sub>3</sub>CH<sub>2</sub>O, J = 7.5 Hz), 8.35 (s, 1H, ArH), 8.77 (s, 1H, lactone-H).

(b) 7,8-Dihydroxy-6-nitro-2-oxo-2*H*-chromene-3-carboxylic acid

- 5        The product from the previous step (a) (0.25 g) was refluxed in 47 % HBr (3 ml) under nitrogen for 6 hours, evaporated and treated with water. Yield: 0.11 g, melting point 287-295° C.

NMR (90 MHz): 8.15 (s, 1H, ArH), 8.74 (s, 1H, lactone-H), 9-12 (br, 3H).

EXAMPLE 3:

- 10        (a) 7,8-Dimethoxy-6-nitro-3-phenyl-chromen-2-one

A mixture of 3,4-dimethoxy-2-hydroxy-5-nitrobenzaldehyde obtained from Preparation example (1a) (0.91 g), phenacetylchloride (1.1 ml) and potassium carbonate (2.0 g) was refluxed in dry acetone (40 ml) overnight. Acetone was evaporated, cold water added and filtered. Yield: 0.69 g.

- 15        NMR (90 MHz): 4.00 (s, 6 H, 2 x CH<sub>3</sub>O), 7.35-7.8 (m, 5 H, Ph), 8.16 (s, 1H, ArH), 8.25 (s, 1H, lactone-H).

(b) 7,8-Dihydroxy-6-nitro-3-phenyl-chromen-2-one

- 20        The product from the previous step (a) (0.6 g) was refluxed in 47 % HBr (6 ml) under nitrogen for 10 hours, poured into ice water and filtered. Yield: 0.38 g, melting point 217-223° C.

NMR (90 MHz): 7.32-7.8 (m, 5H, Ph), 7.98 (s, 1H, ArH), 8.23 (s, 1H, lactone-H).

EXAMPLE 4:

(a) 3-(4-Chloro-phenyl)-7,8-dimethoxy-6-nitro-chromen-2-one

- 25        2-Hydroxy-3,4-dimethoxy-5-nitro-benzaldehyde (2.44 g), 4-chlorophenyl acetic acid (1.82 g), PhOPOCl<sub>2</sub> (1.6 ml) and triethylamine (4.5 ml) in 1,2-dichloroethane (125 ml) were refluxed for 5 hours as described in example 1(a). Yield: 2.40 g.

19

NMR (90 MHz): 4.00 (s, 3H, CH<sub>3</sub>O), 4.02 (s, 3H, CH<sub>3</sub>O), 7.4-7.7 (m, 4H, Ph), 8.16 (s, 1H, ArH), 8.20 (s, 1H, lactone-H).

(b) 3-(4-Chloro-phenyl)-7,8-dihydroxy-6-nitro-chromen-2-one

5 A mixture of the product from the previous step (a) (2g) and 47 % hydrogen bromide (50 ml) was refluxed under nitrogen atmosphere for six hours. After the usual work-up the product was recrystallized from methanol. Yield: 0.85 g.

NMR (90MHz): 7.4-7.7 (m, 4H, Ph), 7.98 (s, 1H, ArH), 8.10 (s, 1H, lactone-H), 8.2-10 (br, 2H, OH).

EXAMPLE 5:

10 (a) 7,8-Dimethoxy-6-nitro-3-o-tolyl-chromen-2-one

A mixture of 2-methylphenyl acetic acid (0.60 g), 2-hydroxy-3,4-dimethoxy-5-nitro-benzaldehyde (0.91 g), PhPOCl<sub>2</sub> (0.60 ml) and triethyl amine (1.7 ml) were reacted in 1,2-dichloroethane as described in example 4. Yield: 0.68 g.

15 NMR (90MHz): 2.22 (s, 3H, CH<sub>3</sub>), 4.03 (s, 3H, CH<sub>3</sub>O), 4.05 (s, 3H, CH<sub>3</sub>O), 7.2-7.4 (m, 4H, Ph), 8.07 (s, 1H, ArH), 8.20 (s, 1H, lactone-H).

(b) 7,8-Dihydroxy-6-nitro-3-o-tolyl-chromen-2-one

The product from the previous step (a) (0.68 g) was refluxed in 47 % HBr (10 ml) for 8 hours under nitrogen and worked up in the usual way, Trituration with boiling ether afforded the product. Yield: 0.24 g, melting point 236-241 °C.

20 NMR (90 MHz): 2.20 (s, 3H, CH<sub>3</sub>), 7.28-7.38 (m, 4H), 7.95 (s, 1H), 7.98 (s, 1H).

EXAMPLE 6:

(a) 7-Hydroxy-6-methoxy-8-nitro-3-phenyl-chromen-2-one

25 To a solution of 7-hydroxy-6-methoxy-3-phenyl-chromen-2-one (1.34 g) in acetone (40 ml) was added nitric acid (0.22 ml in 5 ml of CH<sub>2</sub>Cl<sub>2</sub>) and the resulting mixture stirred for 20 minutes. After evaporation of the solvents the product was recrystallized from ethanol. Yield: 0.85 g.

20

NMR (400 MHz): 3.92 (s, 3H, MeO), 7.52-7.56 (m, 2H, Ph), 7.63 (s, 1H, ArH), 7.67-7.71 (m, 1H, Ph), 7.90-7.92 (m, 2H, Ph), 8.37 (s, 1H, CH=C).

(b) 6,7-Dihydroxy-8-nitro-3-phenyl-chromen-2-one

5 The product from the previous step (a) (0.8 g) was reacted with boron tribromide (1.1 ml) in dichloromethane (25 ml) under nitrogen and in room temperature overnight. The reaction mixture was treated with water, extracted in ethyl acetate and finally recrystallized from ethanol. Yield: 100 mg, melting point 195-210°C.

10 NMR (400 MHz): 7.28 (s, 1H, arH), 7.39-7.42 (m, 3H, Ph), 7.67-7.71 (m, 2H, Ph), 8.23 (s, 1H, CH=C), 10.7 (br, OH).

EXAMPLE 7:

3-Benzoyl-6,7-dihydroxy-5-nitro-chromen-2-one

15 Nitric acid solution (1.6 ml, 2 M in CH<sub>2</sub>Cl<sub>2</sub>) was added to a solution of 3-benzoyl-6,7-dihydroxy-chromen-2-one (0.89 g) in ethyl acetate (50 ml) at -16 - 10°C. The solvent was evaporated and the product run through a silica column with toluene- ethyl acetate acetic acid (8:1:1) as the solvent. The product was crystallized from ether. Yield: 40 mg, melting point 94-96°C.

NMR (400 MHz): 7.02 (s, 1H, ArH), 7.53-7.55 (m, 2H, Ph), 7.69 -7.70 (m, 1H, Ph), 7.87-7.88 (m, 2H, Ph), 8.03 (s, 1H, CH=C), 10-12 (br, 2H, 2 x OH).

20 EXAMPLE 8:

2,3-Dihydroxy-1-nitro-7,8,9,10-tetrahydro-benzo[c]chromen-6-one

25 A solution of 5 M HNO<sub>3</sub> in H<sub>2</sub>SO<sub>4</sub> (1 ml) was added to solution of acetic acid 3-acetoxy-6-oxo-7,8,9,10-tetrahydro-6H-benzo[c]chromen-2-yl ester (1.6 g) in concentrated sulfuric acid (20 ml) and stirred at room temperature for an hour. The mixture was treated with ice (150 g) and filtered. The product was purified by column chromatography on silica (10 % methanol in dichloromethane). Yield: 50 mg, melting point 238-255° C.

NMR (400 MHz): 1.64 (m, 4H), 2.40 (m, 2H), 3.43 (m, 2H), 6.86 (s, 1H), 10.22 (s, 1H), 11.40 (s, 1H).

## EXAMPLE 9:

## (a) 6,7-Dihydroxy-2-imino-2H-chromene-3-carboxylic acid ethyl ester

To a solution of 2,4,5-trihydroxybenzaldehyde (2.94 g) and ethyl cyanoacetate (1.62 g) in DMF (20 ml) were added 20 drops (Pasteur pipette) of piperidine and eight drops of acetic acid and the solution kept at 80 °C under nitrogen for two hours. The product was filtered and washed with DMF and ethyl ether. Yield: 2.00 g.

NMR (300 MHz): 1.31 (t, 3H, J = 7 Hz, CH<sub>2</sub>CH<sub>3</sub>), 4.29 (q, 2H, J = 7 Hz, CH<sub>2</sub>CH<sub>3</sub>), 6.18 (s, 1H, ArH), 6.77 (s, 1H, ArH), 8.12 (s, 1H, CH=C), 8.5 (br, 1H), 8.8 (br, 1H), 9.6 (br, 1H).

## (b) 6,7-Dihydroxy-2-oxo-2H-chromene-3-carboxylic acid

The product from the previous step (a) was refluxed in 1 M hydrochloric acid for four hours, cooled and filtered. Yield: 97 %.

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): 6.82 (s, 1H, ArH), 7.22 (s, 1H, ArH), 8.68 (s, 1H, CH=C-COOH), 9.38 (b, 1H, OH), 10.83 (b, 1H, OH), 12.8 (b, 1H, COOH).

## (c) 6,7-Dihydroxy-5-nitro-2-oxo-2H-chromene-3-carboxylic acid

The product from the previous step (b) (0.22 g) in concentrated sulfuric acid (6 ml) was treated with potassium nitrate (0.10 g) at (-10) °C, poured into ice water and filtered. Yield: 0.21 g, melting point over 300 °C.

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): 6.97 (s, 1H, ArH), 8.26 (s, 1H, CH=C-COOH), 9-13 (br, 3H, 2 x OH + COOH).

## EXAMPLE 10:

## (a) 7-Hydroxy-6-methoxy-2-oxo-2H-chromene-3-carboxylic acid ethyl ester

A mixture of 2,4-dihydroxy-5-methoxybenzaldehyde (1.68 g) and diethyl malonate (1.76 g) in DMF (5 ml) was treated with a few drops of piperidine and acetic acid and kept at 80-95 °C overnight. The reaction mixture was poured into ice water, acidified and filtered. Yield: 2.2.g.

NMR (300 MHz): 1.30 (t, 3H, Et), 3.82 (s, 3H, MeO), 4.26 (q, 2H, Et),

22

6.79 (s, 1H, ArH), 7.44 (s, 1H, ArH), 8.65 (s, 1H, lactone-H), 10.90 (s, 1H, OH).

(b) 7-Hydroxy-6-methoxy-8-nitro-2-oxo-2H-chromene-3-carboxylic acid ethyl ester

5 A solution of 1 M nitric acid in acetic acid (4.1 ml) was added to a suspension of the product from the previous step (a) (1.06 g) in acetic acid (20 ml) at room temperature. The starting suspension was dissolved and the product crystallized out. It was filtered, washed with acetic acid and ethyl ether. Yield: 0.53 g.

10 NMR (300 MHz): 1.30 (t, 3 H, Et), 3.91 (s, 3H, MeO), 4.27 (q, 2H, Et), 7.69 (s, 1H, ArH), 8.72 (s, 1H, lactone-H).

(c) 7-Hydroxy-6-methoxy-8-nitro-2-oxo-2H-chromene-3-carboxylic acid

The product from the previous step (b) (0.5 g) was hydrolyzed by refluxing it in a mixture of 4 M HCl (25 ml) and acetic acid (15 ml) for 30 minutes. The  
15 reaction mixture was agitated in ice bath and filtered. Yield: 0.34 g.

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): 7.35 (s, 1H, ArH), 8.72 (s, 1H, CH=C-COOH), 10.7 (br, 2H, 2 x OH), 12-13 (br, 1H, COOH).

(d) 6,7-Dihydroxy-8-nitro-2-oxo-2H-chromene-3-carboxylic acid

The suspension of the product from the previous step (c) (0.3 g) in  
20 dichloromethane (30 ml) was treated with boron tribromide (1.05 ml) at -25 °C and kept then at room temperature for two days. The reaction mixture was treated with ice water and extracted with ethyl acetate. The product was recrystallized from a mixture of water and 2-propanol. Yield: 0.18 g.

NMR (400 MHz): 7.35 (s, 1H, ArH), 8.73 (s, 1H, lactone-H), 10.7 (br, 1-2H),  
25 12-13 (br, 1H).

#### EXAMPLE 11:

(a) 2-(2,4,5-Trimethoxy-benzylidene)-succinic acid 1-ethyl ester

A solution of 2,4,5-trimethoxybenzaldehyde (19.6 g), diethyl succinate (17.4 g) and potassium tert-butoxide (11.2 g) in ethanol (70 ml) was refluxed for

23

four hours. Ethanol was evaporated, water (400 ml) added and the water solution washed with diethyl ether. The water phase was acidified and the product extracted into ether. Recrystallization from ether yielded an approximately 4:1 mixture of cis-trans isomers. Yield: 22 g.

5 NMR (400 MHz) for the major isomer: 1.24 (t, 3H, Et-CH<sub>3</sub>), 3.65 (s, 2H, CH<sub>2</sub>COOH), 3.65 (s, 3H, Meo), 3.84 (s, 3H, MeO), 4.02 (s, 3H, MeO), 4.19 (q, 2H, Et-CH<sub>2</sub>), 6.75 (s, 1H, ArH), 6.91 (s, 1H, ArH), 7.79 (s, 1H, CH=C), 12.45 (s, 1H, COOH).

(b) (6,7-Dihydroxy-2-oxo-2H-chromen-3-yl)-acetic acid

10 Boron tribromide (3 ml) was added to a solution of 2-(2,4,5-trimethoxybenzylidene)-succinic acid 1-ethyl ester obtained from the previous step (a) (4 g) in dichloromethane (40 ml) at -20°C under nitrogen atmosphere. The reaction was kept on ice for an hour and then at room temperature for three days. After water treatment, filtering and trituration with warm 2-propanol the product was filtered.  
15 Yield: 1.1 g

NMR (400 MHz): 3.39 (s, 2H, CH<sub>2</sub>), 6.75 (s, 1H, ArH), 7.03 (s, 1H, ArH), 7.77 (s, 1H, CH=C), 9.38 (s, 1H, OH), 10.15 (s, 1H, OH), 12.15 (br, 1H, COOH).

(c) (6,7-Dihydroxy-5-nitro-2-oxo-2H-chromen-3-yl)-acetic acid

Potassium nitrate (0.21 g) was added in small doses to a solution of the  
20 product from the previous step (b) (0.46 g) in sulfuric acid at (-18) °C - (-12) °C. The solution was then kept at 0 °C for an hour, poured in ice and filtered. The product was triturated with hot water and filtered at ambient temperature. Yield: 0.15 g, melting point over 350°C.

NMR (400 MHz): 3.51 (s, 2H, CH<sub>2</sub>), 6.94 (s, 1H, ArH), 7.69 (s, 1H, CH=C),  
25 10.5 (br, 1H, OH), 11.6 (br, 1H, OH), 12.47 (br, 1H, COOH).

#### EXAMPLE 12:

(a) 6,7-Dihydroxy-3-(2H-tetrazol-5-yl)-chromen-2-one

A mixture of 2,4,5-trihydroxybenzaldehyde (1.23 g), (2H-tetrazol-5-yl)-acetic acid ethyl ester (1.25 g), piperidine (1.36 g) and acetic acid (0.06 g) in DMF (10 ml)  
30 was stirred at 90°C under nitrogen for two hours. After ice water treatment and



24

acidification to pH 2 the product was filtered and triturated with hot ethanol. Yield: 0.55 g.

NMR (400 MHz): 6.80 (s, 1H), 7.06 (s, 1H, ArH), 8.71 (s, 1H, lactone-H), 10-11 (br, 3H).

5 (b) 6,7-Dihydroxy-5-nitro-3-(1*H*-tetrazol-5-yl)-chromen-2-one

Potassium nitrate (0.14 g) was added to a solution of the product from the previous step (a) (0.4 g) in sulfuric acid (10 ml) at -18 °C - -12 °C. The reaction mixture was poured into ice water and filtered. The product was extracted in hot acetone, concentrated and filtered. Yield: 50 mg, melting point over 315 °C.

10 NMR (400 MHz): 7.08 (s, 1H, ArH), 8.49 (s, 1H, CH=C), 10-12.2 (br, 3H, OH, CN<sub>4</sub>H).

EXAMPLE 13:

(a) (6,7-Dihydroxy-4-methyl-2-oxo-2*H*-chromen-3-yl)-acetic acid ethyl ester

15 To a solution of 1,2,4-trihydroxybenzene (2.5 g) in 75 % sulfuric acid (30 ml) at 5-10 °C was added diethyl 2-acetyl glutarate (4.3 g). The mixture was stirred at ambient temperature overnight poured in ice and filtered. The product (4.2 g) was used as such in the following step (b).

(b) (6,7-Dihydroxy-4-methyl-2-oxo-2*H*-chromen-3-yl)-acetic acid

20 A solution of (6,7-dihydroxy-4-methyl-2-oxo-2*H*-chromen-3-yl)-acetic acid ethyl ester obtained from the previous step (a) (4.2 g) in acetic acid (20 ml) and 6 N hydrochloric acid (50 ml) was refluxed for two hours, cooled and filtered. Yield: 2.3 g.

25 NMR (400 MHz): 2.28 (s, 3H, CH<sub>3</sub>), 3.62 (s, 2H, CH<sub>2</sub>COOH), 6.74 (s, 1H, ArH), 7.07 (s, 1H, ArH), 9.34 (s, 1H, OH), 10.19 (s, 1H, OH).

(c) (6,7-Dihydroxy-4-methyl-5-nitro-2-oxo-2*H*-chromen-3-yl)-acetic acid

To a solution of the product from the previous step (b) (2 g) in concentrated sulfuric acid (10 ml) at -15 °C was added potassium nitrate (1.02 g). The mixture


25

was kept at 0 °C for an hour, treated with ice water and filtered. Yield: 0.41 g, melting point over 350°C.

NMR(400 MHz): 2.49 (s, 3H, CH<sub>3</sub>), 3.57 (s, 2H, CH<sub>2</sub>), 6.92 (s, 1H, ArH), 10.3 (br, 1H, OH), 11.6 (br, 1H, OH).

## CLAIMS:

1. A compound of formula I



wherein the two OH- substituents in the phenyl moiety are in a position ortho to one another and R1 in a position ortho to one of the hydroxy groups;

10 X is O or NR<sub>7</sub>; wherein R<sub>7</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or -(C<sub>1</sub>-C<sub>6</sub>)alkyl-COOH;  
R<sub>1</sub> is NO<sub>2</sub>, CN, CHO, CF<sub>3</sub> or (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-;

R<sub>2</sub> and R<sub>3</sub> are each selected independently from H, OH, halogen, NO<sub>2</sub>, SH, NH<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, OH-(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, SO<sub>2</sub>R<sub>10</sub>, -CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl, -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH and -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>; wherein


m is 0 or 1;

n is 0 or 1;

Y is -CO- or -CHOH-;

20 B is (C<sub>1</sub>-C<sub>6</sub>)alkylene or (C<sub>2</sub>-C<sub>6</sub>)alkenylene;

R<sub>8</sub> is phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclyl with one to four heteroatoms each selected independently from N, O and S, wherein the said phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclyl is optionally substituted with one to five substituents R<sub>9</sub> each selected independently from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub>, SH, NH<sub>2</sub>, CN, CHO, =O, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, CO-NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino-CO-, NHOH, CONHOH or SO<sub>2</sub>R<sub>10</sub>;  
30 or R<sub>8</sub> is



or R<sub>2</sub> and R<sub>3</sub> form together -(CH<sub>2</sub>)<sub>r</sub>; wherein r is 3, 4 or 5;


- 5 R<sub>4</sub> and R<sub>5</sub> are independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or R<sub>4</sub> and R<sub>5</sub> form together =O, =S or =NR<sub>11</sub>, wherein R<sub>11</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
 R<sub>6</sub> is H, NO<sub>2</sub>, CN, CHO, halogen, CF<sub>3</sub> or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and  
 R<sub>10</sub> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>, OH or mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino;

- 10 with the provisos that (a) when X is O, R<sub>2</sub> is methyl, R<sub>3</sub> is H, R<sub>4</sub> and R<sub>5</sub> form together =O, R<sub>6</sub> is H and the two hydroxy substituents are at 7- and 8-positions, then R<sub>1</sub> is not CHO, (b) when X is O, R<sub>2</sub> is H or methyl, R<sub>3</sub> is H, R<sub>4</sub> and R<sub>5</sub> form together =O, R<sub>6</sub> is H and the two hydroxy substituents are at 6- and 7-positions, then R<sub>1</sub> is not 8-CO-CH<sub>3</sub>, (c) when X is O, R<sub>2</sub> and R<sub>3</sub> are H, R<sub>4</sub> and R<sub>5</sub> are both  
 15 methyl, R<sub>6</sub> is H and the two hydroxy substituents are at 7- and 8-positions, then R<sub>1</sub> is not -CO-CH<sub>3</sub>,

or pharmaceutically acceptable salts or pharmaceutically acceptable esters thereof.

2. A compound according to claim 1, which is a compound of formula Ia,


20



wherein X, R<sub>1</sub> to R<sub>5</sub> are as defined in claim 1.

28

3. A compound according to claim 1, which is a compound of formula Ib,



wherein X, R<sub>1</sub> to R<sub>5</sub> are as defined in claim 1.

4. A compound according to any one of claims 1 to 3, wherein X is O.

5. A compound according to any one of claims 1 to 4, wherein R<sub>1</sub> is NO<sub>2</sub>, CN or CF<sub>3</sub>.

6. A compound according to any one of claims 1 to 4, wherein R<sub>4</sub> and R<sub>5</sub> form together =O.

7. A compound according to any one of claims 1 to 5, wherein R<sub>3</sub> is

10 -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH or -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>.

8. A compound according to any one of claims 1 to 7, wherein R<sub>8</sub> is phenyl or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, each unsubstituted or substituted with one or two substituent(s) R<sub>9</sub> each selected independently from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub> and (C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>8</sub> is 5-tetrazolyl.


15 9. A compound according to any one of claims 1 to 8 for use as a pharmaceutical.

10. A pharmaceutical composition comprising at least one compound according to any one of claims 1 to 8 and a pharmaceutically acceptable excipient.

11. A pharmaceutical composition according to claim 10, wherein the two hydroxy substituents in the phenyl moiety are in 6- and 7-positions.

12. A pharmaceutical composition according to claim 10, wherein the two hydroxy substituents in the phenyl moiety are in 7- and 8-positions.

13. Use of a compound of formula I',



29

wherein the two OH- substituents in the phenyl moiety are in a position ortho to one another and R<sub>1</sub> in a position ortho to one of the hydroxy groups;

X is O or NR<sub>7</sub>; wherein R<sub>7</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or -(C<sub>1</sub>-C<sub>6</sub>)alkyl-COOH;

5 R<sub>1</sub> is NO<sub>2</sub>, CN, CHO, CF<sub>3</sub> or (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-;

R<sub>2</sub> and R<sub>3</sub> are each selected independently from H, OH, halogen, NO<sub>2</sub>, SH, NH<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, OH-(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, SO<sub>2</sub>R<sub>10</sub>, -CO-(C<sub>1</sub>-C<sub>9</sub>)alkyl, -(Y)<sub>n</sub>-(B)<sub>m</sub>-COOH and -(Y)<sub>n</sub>-(B)<sub>m</sub>-R<sub>8</sub>; wherein

m is 0 or 1;

n is 0 or 1;


Y is -CO- or -CHOH-;

B is (C<sub>1</sub>-C<sub>6</sub>)alkylene or (C<sub>2</sub>-C<sub>6</sub>)alkenylene;

15

R<sub>8</sub> is phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclyl with one to four heteroatoms each selected independently from N, O and S, wherein the said phenyl, naphthyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or 5- or 6-membered heterocyclyl is optionally substituted with one to five substituents R<sub>9</sub> each selected independently from OH, halogen, COOH, 5-tetrazolyl, NO<sub>2</sub>, SH, NH<sub>2</sub>, CN, CHO, =O, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo-(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, CO-NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino-CO-, NHOH, CONHOH or SO<sub>2</sub>R<sub>10</sub>;

or R<sub>8</sub> is



25

or R<sub>2</sub> and R<sub>3</sub> form together -(CH<sub>2</sub>)<sub>r</sub>; wherein r is 3, 4 or 5;

R<sub>4</sub> and R<sub>5</sub> are independently H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; or R<sub>4</sub> and R<sub>5</sub> form together =O, =S or =NR<sub>11</sub>, wherein R<sub>11</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

30 R<sub>6</sub> is H, NO<sub>2</sub>, CN, CHO, halogen, CF<sub>3</sub> or (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

R<sub>10</sub> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>, OH or mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino;

or pharmaceutically acceptable salts or pharmaceutically acceptable esters thereof, for the manufacture of a medicament as a COMT inhibiting agent.


30

14. A use according to claim 13, wherein the two hydroxy substituents in the phenyl moiety are in 6- and 7-positions.

15. A use according to claim 13, wherein the two hydroxy substituents in the phenyl moiety are in 7- and 8-positions.

5 16. A use according to claim 13, wherein a compound of formula I' is a compound of formula I as defined in any one of claims 1, 2, 3, 4, 5, 6, 7 and/or 8.

17. A method for the treatment of diseases or conditions where inhibitors of COMT enzyme are indicated to be useful, said method comprising administering to a mammal in need of such treatment an effective amount of a compound of  
10 formula I',



wherein the two OH- substituents in the phenyl moiety are in a position ortho to one another and R<sub>1</sub> in a position ortho to one of the hydroxy groups;  
15 and X, R<sub>1</sub> to R<sub>6</sub> are as defined in claim 9.

18. A method according to claim 17, wherein the two hydroxy substituents in the phenyl moiety are in 6- and 7-positions.

19. A method according to claim 17, wherein the two hydroxy substituents in the phenyl moiety are in 7- and 8-positions.

20 20. A method according to claim 17, wherein a compound of formula I' is a compound of formula I as defined in any one of claims 1, 2, 3, 4, 5, 6, 7 and/or 8.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/FI 01/00613

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D311/04 C07D215/20 A61K31/47 A61P25/16 A61K31/435

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K C07C A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

PAJ, EP0-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PIA LAUTALA ET AL: "Molecular Mechanisms Controlling the Rate and Specificity of Catechol O-Methylation by Human Soluble Catechol O-Methyltransferase" MOLECULAR PHARMACOLOGY, vol. 69, no. 2, 2001, pages 393-402, XP002901963 DATABASE: CAPLUS, RN 305-01-1 figure 2 --- -/--	1-20

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 October 2001

Date of mailing of the international search report

07 12 2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Farieta Fernando



## INTERNATIONAL SEARCH REPORT

 tional Application No  
 PCT/FI 01/00613

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ERDEM YESILADA ET AL: "In vitro inhibitory effects of daphne oleoides ssp. oleoides on inflammatory cytokines and activity-guided isolation of active constituents" CYTOKINE, vol. 13, no. 6, March 2001 (2001-03), pages 359-364, XP002901954 DATABASE: CAPLUS, RN 486-35-1P; compound 9 figure 1	1-20
X	--- PATENT ABSTRACTS OF JAPAN vol. 1995, no. 06, 31 July 1995 (1995-07-31) & JP 07 076583 A (MORINAGA MILK IND CO LTD), 20 March 1995 (1995-03-20) abstract	1-20
A	--- US 5 446 194 A (BAECKSTROEM REIJO J ET AL) 29 August 1995 (1995-08-29) claims 1-4	1-20
A	--- US 5 389 653 A (BERNAUER KARL ET AL) 14 February 1995 (1995-02-14) claims 1-23	1-20
A	--- SEPPO KAAKKOLA: "Clinical Pharmacology, Therapeutic Use and Potential of COMT Inhibitors in Parkinson's Disease" DRUGS, vol. 59, no. 6, June 2000 (2000-06), pages 1233-1250, XP002901955 the whole document	1-20
A	--- GIUSEPPE MECO ET AL: "Can COMT Inhibitors Improve Cognitive Functions in Patients with Parkinson's Disease" CNS DRUGS, vol. 13, no. 4, April 2000 (2000-04), pages 227-232, XP002901956 the whole document	1-20
A	--- CHERYL WATERS MD FRCP(C): "Catechol-O-Methyltransferase (COMT) Inhibitors in Parkinson's Disease" JAGS, vol. 48, 2000, pages 692-698, XP002901957 the whole document -----	1-20

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/FI 01/00613

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
JP 07076583	A	20-03-1995	JP	3166094 B2	14-05-2001
			AU	673916 B2	28-11-1996
			AU	6438094 A	08-11-1994
			CA	2138021 A1	27-10-1994
			DE	69425210 D1	17-08-2000
			DE	69425210 T2	01-03-2001
			EP	0645382 A1	29-03-1995
			WO	9424119 A1	27-10-1994
			US	5574062 A	12-11-1996
-----					
US 5446194	A	29-08-1995	US	5283352 A	01-02-1994
			AT	401053 B	25-06-1996
			AT	312987 A	15-10-1995
			AU	621036 B2	05-03-1992
			AU	8187987 A	02-06-1988
			BE	1003279 A5	18-02-1992
			BG	60407 B2	28-02-1995
			CA	1289078 A1	17-09-1991
			CA	1334967 A1	28-03-1995
			CH	685436 A5	14-07-1995
			CN	87108011 A ,B	08-06-1988
			CS	276263 B6	13-05-1992
			CS	8808439 A3	15-01-1992
			CS	277018 B6	18-11-1992
			CS	8808440 A3	13-05-1992
			DD	281375 A5	08-08-1990
			DE	3740383 A1	01-06-1988
			DK	623087 A	29-05-1988
			EG	18338 A	30-09-1992
			ES	2008359 A6	16-07-1989
			FI	875229 A ,B,	29-05-1988
			FR	2607493 A1	03-06-1988
			GB	2200109 A ,B	27-07-1988
			GR	871817 A1	24-03-1988
			HK	75594 A	12-08-1994
			HR	921250 A1	30-06-1998
			HU	45473 A2	28-07-1988
			IE	60320 B	29-06-1994
			IT	1225762 B	26-11-1990
			JP	2735834 B2	02-04-1998
			JP	63150237 A	22-06-1988
			JP	2096417 C	02-10-1996
			JP	8005781 B	24-01-1996
			JP	63170311 A	14-07-1988
			KR	9604771 B1	13-04-1996
			LT	915 A ,B	27-03-1995
			LU	87050 A1	03-05-1988
			LV	10236 A ,B	20-10-1994
			MT	1012 A	18-10-1988
			MX	9524 A	01-12-1993
			NL	8702857 A	16-06-1988
			NO	874966 A ,B,	30-05-1988
			NZ	222729 A	26-04-1990
			PH	26145 A	18-03-1992
			PL	269091 A1	08-12-1988
			PT	86236 A ,B	01-12-1987
			SE	503434 C2	17-06-1996
			SE	8704751 A	29-05-1988

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/FI 01/00613

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 5446194	A		RU	2014319 C1	15-06-1994
			SU	1729291 A3	23-04-1992
-----					
US 5389653	A	14-02-1995	US	5236952 A	17-08-1993
			US	5476875 A	19-12-1995
			US	5633371 A	27-05-1997
			US	5705703 A	06-01-1998
			AT	90072 T	15-06-1993
			AU	603788 B2	29-11-1990
			AU	6976487 A	17-09-1987
			CA	1302418 A1	02-06-1992
			CS	9104022 A3	16-12-1992
			DE	3786026 D1	08-07-1993
			DK	82087 A	12-09-1987
			EP	0237929 A1	23-09-1987
			ES	2056792 T3	16-10-1994
			FI	871041 A ,B,	12-09-1987
			HK	165896 A	13-09-1996
			HU	43804 A2	28-12-1987
			IE	61316 B	02-11-1994
			IL	81791 A	01-12-1992
			JP	2006568 C	11-01-1996
			JP	7042254 B	10-05-1995
			JP	62240649 A	21-10-1987
			KR	9301054 B1	13-02-1993
			KR	9301337 B1	26-02-1993
			LU	90219 A9	28-05-1998
			LV	5742 A4	20-10-1996
			MC	1807 A	22-12-1987
			NO	870984 A ,B,	14-09-1987
			NZ	219496 A	21-12-1990
			PT	84449 A ,B	01-04-1987
			ZA	8701564 A	28-10-1987
			MX	9203634 A1	01-07-1992
-----					